

AD-A193 481	MONOTONE RECURRENCE RELATIONS THEIR BIRKHOFF ORBITS AND TOPOLOGICAL ENTRO (U) WISCONSIN UNIV-MADISON CENTER FOR MATHEMATICAL SCIENCES 5 B ANGENENT FEB 88	1/1
UNCLASSIFIED	CMS-TSR-88-25 DAAL03-87-K-0043	F/G 12/2 NL

MONOTONE RECURRENCE RELATIONS THEIR BIRKHOFF ORBITS AND
TOPOLOGICAL ENTRO (U) WISCONSIN UNIV-MADISON CENTER
FOR MATHEMATICAL SCIENCES S B ANGENENT FEB 88
CMS-TSR-88-25 DAAL03-87-K-0043 F/G 12/2


1/1

UNCLASSIFIED

F/G 12/2

NL

111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929



AD-A193 481

CMS Technical Summary Report #88-25

MONOTONE RECURRENCE RELATIONS,
THEIR BIRKHOFF ORBITS AND
TOPOLOGICAL ENTROPY

Sigurd B. Angenent

UNIVERSITY
OF WISCONSIN



CENTER FOR THE
MATHEMATICAL
SCIENCES

Center for the Mathematical Sciences
University of Wisconsin—Madison
610 Walnut Street
Madison, Wisconsin 53705


February 1988

(Received November 20, 1987)

Approved for public release
Distribution unlimited

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709



88 3 23 03 6

UNIVERSITY OF WISCONSIN - MADISON
CENTER FOR THE MATHEMATICAL SCIENCES

MONOTONE RECURRENCE RELATIONS,
THEIR BIRKHOFF ORBITS AND TOPOLOGICAL ENTROPY

Sigurd B. Angenent*

Technical Summary Report #88-25

February 1988

ABSTRACT

Sub 1
A generalization of the class of monotone twistmaps to maps of $S^1 \times \mathbb{R}^N$ is proposed. The existence of Birkhoff orbits is studied, and a criterion for positive topological entropy is given. These results are then specialized to the case of monotone twist maps. Finally it is shown that there is a large class of symplectic maps to which the foregoing discussion applies. (Keywords: dynamical systems, theorems).

AMS (MOS) Subject Classification: 58F12, 58F99

Key Words: Twist map, higher dimensional annulus, Birkhoff orbit, topological entropy

This research was partially supported by the United States Army under Contract No. DAAL03-87-K-0043, by a NATO Science Fellowship, and by the Netherlands Organization for the Advancement of Pure Research (Z.W.O.).

CONTENTS

1. Introduction
2. Twist maps and recurrence relations
3. Birkhoff orbits and the space X
4. Subsolutions and supersolutions
5. A generalization of a theorem of Hall's
6. Quasiperiodic orbits with prescribed rotation number
7. A criterion for positive entropy
8. Twist maps with zero entropy
9. Existence of Birkhoff sub- and super-solutions in general
10. The symplectic case

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



MONTONE RECURRENCE RELATIONS,
THEIR BIRKHOFF ORBITS AND TOPOLOGICAL ENTROPY

Sigurd Angenent*

§1. Introduction.

In this paper we shall study a class of dynamical systems which generalizes the class of "monotone twist maps" of two dimensional annuli, and also the class of "degree one circle maps", both of which have been studied extensively in the last decade or two.

The maps which we shall deal with are continuous maps φ of the generalized annulus $S^1 \times \mathbb{R}^N$ ($N \geq 0$) into itself. The defining condition of these maps is that they come from solving a recurrence relation

$$(1.1) \quad \Delta(x_{k-l}, \dots, x_{k+m}) = 0 \quad (\forall k \in \mathbb{Z})$$

in the first coordinate of a general point in $S^1 \times \mathbb{R}^N$. This is explained more precisely in section two.

Our main concern is with the construction of Birkhoff orbits of φ (defined in section three), and a criterion for positivity of the topological entropy, $h_{\text{top}}(\varphi)$, of the map φ .

The main tool which we use is presented in section four, in the form of theorem 4.2 and two afterthoughts. This theorem allows us to construct solutions of the recurrence equation with prescribed qualitative properties. The method of construction is a discrete analog of Perron's method of solving the Dirichlet problem i.e. his method of constructing harmonic functions with prescribed boundary values from subharmonic functions and "barrier functions" (or super harmonic functions).

This research was partially supported by the United States Army under Contract No. DAAL03-87-K-0043, by a NATO Science Fellowship and by the Netherlands Organization for the Advancement of Pure Research (Z.W.O.).

As a first application of this method we show in section five how the existence of certain kinds of orbits of φ implies the existence of a Birkhoff orbit, thereby generalizing a result of G. R. Hall ([Ha 1]).

In section six we formulate hypothesis 6.1, which roughly speaking, asserts the existence of two solutions of (1.1), say $x_n^{(1)}$ and $x_n^{(2)}$, which "exchange rotation numbers" in the sense that

$$\lim_{n \rightarrow \infty} \frac{x_n^{(1)}}{n} \geq \omega_1 \geq \lim_{n \rightarrow -\infty} \frac{x_n^{(2)}}{n}$$

and

$$\lim_{n \rightarrow \infty} \frac{x_n^{(2)}}{n} \leq \omega_0 \leq \lim_{n \rightarrow -\infty} \frac{x_n^{(1)}}{n}$$

hold for some $\omega_0 < \omega_1$. It is shown that this hypothesis implies the existence of Birkhoff orbits with any prescribed rotation number ω in the interval $[\omega_0, \omega_1]$.

A more complicated construction, which is carried out in section seven, shows that the hypothesis (6.1) implies that $h_{\text{top}}(\varphi) > 0$. In fact we show that there is a compact set $K \subset S^1 \times \mathbb{R}^N$ such that some iterate of φ leaves K invariant, and has a Bernoulli shift as a factor, when restricted to K . Here we partially generalize another result of G. R. Hall, who in turn was concerned with giving a topological version of a "shadowing theorem" of J. N. Mather's (see [Ha 2] and [Ma 2] where Mather's result is announced). A consequence of this construction is that the number of periodic orbits with period $\leq N$ grows exponentially with N . Moreover, our method of proving existence of these orbits is constructive. Given $x^{(1)}$ and $x^{(2)}$ a computer program could be written which constructs a great number of periodic orbits.

In section eight we specialize our results to the two dimensional case, and ask which properties a map (i.e. a monotone twist map) with zero

topological entropy must have. The first result in this direction is that, if $h_{\text{top}}(\varphi) = 0$, then any orbit must have a forward and backward rotation number. Similar results, under different hypotheses, have been obtained by M. Handel. The other result is that, if $h_{\text{top}}(\varphi) = 0$, then any periodic orbit "of type (p,q) " with $\gcd(p,q) = 1$ must be a Birkhoff orbit. This result is originally due to P. Boyland [Bo]. As far as I am aware, both Handel and Boyland rely on Thurston's classification of surface diffeomorphisms. By contrast, our approach is self contained, and as we have pointed out before, constructive, in a certain sense.

Throughout section eight we could have weakened the hypothesis " $h_{\text{top}}(\varphi) = 0$ " to "there is no invariant set $K \subset S^1 \times \mathbb{R}^1$ such that $\varphi|_K$ has a subshift of finite type as a factor. However, a theorem of Katok ([Ka 2]) shows that this is not very much of a weakening. Indeed, if the map φ is $C^{1+\epsilon}$, then his theorem says that both conditions are equivalent.

The next section, the ninth, deals with the question of existence of Birkhoff orbits in general, i.e. without further hypotheses on the map φ . We obtain the existence of Birkhoff orbits with prescribed rotation number ω , not for the original map φ , but for a "translated map" which comes from the recurrence relation $\Delta(x_{k-l}, \dots, x_{k+m}) = \lambda$ for some λ . In general this λ will depend on ω , and will not be zero. From this result we derive a sufficient condition of the existence of at least one Birkhoff orbit of the original map φ . The condition turns out to be (trivially) necessary as well. We conclude this section by briefly specializing the results to the one dimensional (i.e. degree-one circle maps) and the two dimensional (twist map) case.

Finally, in the last section, by way of example, we indicate a number of symplectic maps φ which belong to the class of maps which was studied in

this paper. In particular we show that there is a very simple mechanical model, whose steady states are described by such maps.

§2. Twist maps and recurrence relations.

Let A be the cylinder $(\mathbb{R}/\mathbb{Z}) \times \mathbb{R}$, and let $\varphi : A \rightarrow A$ be an orientation preserving monotone twist homeomorphism. On the universal cover \mathbb{R}^2 of A a lift of φ will be given by

$$F(x,y) = (f(x,y), g(x,y)) ,$$

where f, g are periodic in the sense that

$$(2.1) \quad \begin{aligned} f(x+1, y) &= f(x,y) + 1 \\ g(x+1, y) &= g(x,y) \end{aligned}$$

(i.e. x is the angle coordinate). The monotone twist condition on φ says that the function $f(x,y)$ is strictly increasing in y .

In addition we shall assume that φ satisfies the infinite twist condition:

$$\lim_{y \rightarrow \pm\infty} f(x,y) = \pm\infty .$$

Combined, these two conditions imply that for any pair $x, \bar{x} \in \mathbb{R}$ there is a unique solution $Y(x, \bar{x})$ of the equation

$$f(x, Y(x, \bar{x})) = \bar{x} .$$

This solution is a strictly increasing function of \bar{x} . It is continuous, and satisfies

$$Y(x, \bar{x}) = Y(x+1, \bar{x}+1)$$

for all $x, \bar{x} \in \mathbb{R}$.

From $Y(x, \bar{x})$ we construct another function:

$$\bar{Y}(x, \bar{x}) = g(x, Y(x, \bar{x})) .$$

This function is also continuous, and periodic in the sense that $\bar{Y}(x, \bar{x}) \equiv \bar{Y}(x+1, \bar{x}+1)$. We shall now show that $Y(x, \bar{x})$ is a strictly decreasing function of x .

For fixed x_0 the image of the line $x = x_0$ under the map F is given by the graph of the function $G_0(\bar{x}) = g(x_0, Y(x_0, \bar{x}))$. This graph divides the plane into two parts, one above it, and one below it. If $x_1 > x_0$ is given, then the image of the line $x = x_1$ must lie in one of these two parts. Thus we either have $g(x_0, Y(x_0, \bar{x})) > g(x_1, Y(x_1, \bar{x}))$ for all $\bar{x} \in \mathbb{R}$, or we have the reverse inequality for all $\bar{x} \in \mathbb{R}$. Since φ and F are orientation preserving the latter cannot happen. Therefore $x_0 < x_1$ implies that $\bar{Y}(x_0, \bar{x}) > \bar{Y}(x_1, \bar{x})$.

The construction of Y also shows that

$$\lim_{x \rightarrow +\infty} Y(x, \bar{x}) = +\infty$$

and similarly $\lim_{x \rightarrow -\infty} Y(x, \bar{x}) = -\infty$.

Consider a sequence of points (x_k, y_k) ($k \in \mathbb{Z}$) in the plane. They are the orbit of a point under the map F if and only if they satisfy

$$x_{k+1} = f(x_k, y_k)$$

$$y_k = g(x_{k-1}, y_{k-1})$$

for all $k \in \mathbb{Z}$. The first equation is equivalent to $y_k = Y(x_k, x_{k+1})$ for all k . If we substitute this in the second equation we get the following equivalent set of equations:

$$y_k = Y(x_k, x_{k+1})$$

$$y_k = \bar{Y}(x_{k-1}, x_k) .$$

We see that a sequence of points (x_k, y_k) is an orbit of F if and only if the x coordinates satisfy

$$(2.2) \quad \Delta(x_{k-1}, x_k, x_{k+1}) = 0 \quad (\forall k \in \mathbb{Z})$$

and the y_k are given by $y_k = Y(x_k, x_{k+1})$. Here we have written

$$(2.3) \quad \Delta(x_{-1}, x_0, x_1) = Y(x_0, x_1) - \bar{Y}(x_{-1}, x_0) .$$

So instead of studying orbits of the map F one may as well study solutions of (2.2).

The function Δ which we have just introduced satisfies the following hypotheses. It is continuous, it is monotone increasing in both x_{-1} and x_{+1} , and

$$\lim_{x_{-1} \rightarrow +\infty} \Delta(x_{-1}, x_0, x_1) = \lim_{x_1 \rightarrow +\infty} \Delta(x_{-1}, x_0, x_1) = +\infty .$$

Finally, it is periodic in the sense that

$$\Delta(x_{-1}, x_0, x_1) \equiv \Delta(x_{-1}+1, x_0+1, x_1+1)$$

holds.

Motivated by this example we shall consider **monotone recurrence relations of type (ℓ, m)** for $\ell, m > 1$. We define such a relation to be one of the form

$$(2.4) \quad \Delta(x_{k-\ell}, \dots, x_{k+m}) = 0 \quad (k \in \mathbb{Z})$$

where Δ is a continuous function of $\ell + m + 1$ variables which satisfies the following conditions:

(2.5): a monotonicity $\Delta(x_{-\ell}, \dots, x_m)$ is a nondecreasing function of all the x_j except possibly x_0 . Moreover, it is strictly increasing in the variables $x_{-\ell}$ and x_{+m} .

b periodicity $\Delta(x_{-\ell}, \dots, x_m) \equiv \Delta(x_{-\ell}+1, \dots, x_m+1)$.

c coerciveness $\lim_{x_{-\ell} \rightarrow +\infty} \Delta(x_{-\ell}, \dots, x_m) = +\infty$ and $\lim_{x_m \rightarrow +\infty} \Delta(x_{-\ell}, \dots, x_m) = +\infty$.

If $(x_{k-l}, \dots, x_{k+m-1})$ is given then conditions a and c imply that we can solve (2.4) for x_{k+m} . In this way we have defined a continuous map F_Δ from \mathbb{R}^{l+m} to \mathbb{R}^{l+m} , given by

$$F_\Delta(x_{k-l}, \dots, x_{k+m-1}) = (x_{k-l+1}, \dots, x_{k+m})$$

The conditions a and c also imply that we can solve (2.4) for x_{k-l} if the other variables are given. Thus the map F_Δ is a homeomorphism of \mathbb{R}^{l+m} onto itself.

Throughout the paper we shall keep one fixed \mathbb{Z} action on \mathbb{R}^{l+m} in mind. It is given by

$$(x_{-l}, \dots, x_{m-1}) + i = (x_{-l} + i, \dots, x_{m-1} + i)$$

for $i \in \mathbb{Z}$. The quotient $\mathbb{R}^{l+m}/\mathbb{Z}$ is homeomorphic to $S^1 \times \mathbb{R}^{l+m-1}$. We shall call it A^{l+m} , since it generalizes the annulus $(\mathbb{R}/\mathbb{Z}) \times \mathbb{R}$ which we started with.

The periodicity condition b implies that F_Δ is equivariant with respect to the \mathbb{Z} -action on \mathbb{R}^{l+m} , i.e. $F_\Delta(x+i) = F_\Delta(x) + i$. Hence it defines a homeomorphism on the quotient A^{l+m} , which we shall call φ_Δ . This class of homeomorphisms is our generalization of the class of monotone twist maps of the annulus.

Our assumption that $l > 1$ is not a necessary assumption. If we allow $l = 0$ then we can still define the maps F_Δ and φ_Δ as above. The only difference is that they need not be invertible anymore. Except for this all the results which we shall derive in the following sections remain true. In particular they apply to degree-one maps of the circle. If $h : \mathbb{R} \rightarrow \mathbb{R}$ satisfies $h(x+1) = h(x) + 1$ then the solutions of (2.4) where we have taken

$$\Delta(x_0, x_1) = x_1 - h(x_0)$$

are exactly the orbits of the map $h : \mathbb{R} \rightarrow \mathbb{R}$. Our map φ_Δ is the map which h induces on the circle \mathbb{R}/\mathbb{Z} .

§3. Birkhoff orbits, and the space X .

Let X denote the space of bi-infinite sequences of real numbers, i.e. $X = \mathbb{R}^{\mathbb{Z}}$. We equip X with the product topology.

The space X is also partially ordered by:

$$x \leq y \stackrel{\text{def}}{\iff} x_k \leq y_k \text{ for all integers } k$$

where $x, y \in X$. The following notation is sometimes used:

$$x < y \stackrel{\text{def}}{\iff} x < y \text{ and } x \neq y,$$

$$x \ll y \stackrel{\text{def}}{\iff} x_k < y_k \text{ for all } k \in \mathbb{Z}.$$

Given a pair of sequences $x, y \in X$ such that $x < y$ we define the **order interval**

$$[x, y] = \{z \in X : x \leq z \leq y\}.$$

Any order interval is homeomorphic to the product $[0, 1]^{\mathbb{Z}}$, which is a compact metrizable space.

On X we have a $\mathbb{Z} \times \mathbb{Z}$ action, τ , given by

$$\tau_{m,n}(x)_i = x_{i-m} + n.$$

This action is compatible with both the topology and the partial ordering we have on X .

A sequence $x \in X$ will be called a **Birkhoff-sequence** if for any pair of integers (m, n) one either has $\tau_{m,n}(x) > x$ or $\tau_{m,n}(x) < x$. We shall denote the set of all Birkhoff sequences by B .

An equivalent definition of a Birkhoff sequence is the following. A sequence $x \in X$ is a Birkhoff sequence if and only if for any $i, j, k \in \mathbb{Z}$ one has

$$x_i < x_j + k \iff x_{i+1} < x_{j+1} + k.$$

Thus, if $\{x_i\}$ is the sequence of x -coordinates of an orbit of a monotone

twist map, then x is a Birkhoff sequence if and only if the map preserves the order on the orbit. This is the usual definition of a "Birkhoff-orbit" (see [Ka 1]).

It is known that if $x \in B$ is a Birkhoff sequence, then x has a rotation number, i.e. the limit

$$\lim_{k \rightarrow +\infty} \frac{x_k}{k} = \omega \stackrel{\text{def}}{=} \omega(x)$$

exists. In fact one has the following inequalities.

$$(3.1) \quad x_0 + \lfloor \omega k \rfloor \leq x_k < x_0 + \lfloor \omega k \rfloor + 1 \quad (k \in \mathbb{Z})$$

where $\lfloor \omega k \rfloor$ is the largest integer below ωk .

The set of Birkhoff sequences, B , which is the intersection of the sets $\{x \in X : \tau_{m,n}(x) > x \text{ or } \tau_{m,n}(x) < x\}$, is a closed subset of X .

The whole set B is not compact, but the inequalities (3.1) imply that for any two constants $\alpha, \beta > 0$ the closed set

$$\{x \in B \mid |x_0| < \alpha \text{ and } |\omega(x)| < \beta\}$$

is contained in an order interval, and therefore compact.

We shall occasionally say something about periodic orbits "of type (p,q) ".

If the map φ_Δ of $\mathbb{R}^{l+m}/\mathbb{Z}$ has a periodic point P with period q , then some lift $(x_{-l}, \dots, x_{m-1}) \in \mathbb{R}^{l+m}$ of P will almost be periodic in the sense that the corresponding sequence $\{x_k, k \in \mathbb{Z}\} \in X$ satisfies

$$(3.2) \quad x_{k+q} = x_k + p \quad (k \in \mathbb{Z})$$

for some $p \in \mathbb{Z}$. Another way of saying this is $x = \tau_{p,q}(x)$.

By definition any sequence $x \in X$ will be said to be periodic of type (p,q) if it satisfies (3.2).

If $x \in X$ is periodic of type (p,q) and if x is also a Birkhoff orbit or sequence then x is periodic of type (p_0, q_0) where $p = l \cdot p_0$,

$q = l \cdot q_0$ and $l = \gcd(p, q)$.

Indeed, $x \in B$ implies that we have $\tau_{p_0, q_0}(x) > x$ or $\tau_{p_0, q_0}(x) < x$.

Suppose the first holds, and for some $i \in \mathbb{Z}$ we have

$$\tau_{p_0, q_0}(x)_i > x_i, \text{ i.e. } x_{i-q_0} + p_0 > x_i.$$

Then $\tau_{p_0, q_0}(x) > x$ implies that for all $k \in \mathbb{Z}$

$$x_{i-kq_0} + kp_0 > x_{i-(k-1)q_0} + (k-1)p_0$$

so that $x_i = x_{i-lq_0} + lp_0 > x_i$, a clear contradiction. If we didn't have

$\tau_{p_0, q_0}(x) > x$ but $\tau_{p_0, q_0}(x) < x$ instead then the same argument would show that we still have $\tau_{p_0, q_0}(x) = x$.

§4. Substitutions and supersolutions.

Let $\Delta \in C(\mathbb{R}^{l+m+1})$ be a monotone recurrence relation of type (l, m) as we defined them in section 2.

A sequence $x \in X$ is called a **subsolution** for Δ iff

$$(4.1) \quad \Delta(x_{j-l}, \dots, x_{j+m}) > 0 \quad (\forall j \in \mathbb{Z})$$

holds. Similarly, a **supersolution** is a sequence $x \in X$ such that

$$(4.1') \quad \Delta(x_{j-l}, \dots, x_{j+m}) < 0 \quad (\forall j \in \mathbb{Z})$$

holds.

The next lemma states some of the elementary properties of sub- and super-solutions.

Lemma 4.1. (a) The set of subsolutions for Δ is closed in X .

The same holds for supersolutions.

(b) If $\{x^{(\alpha)}\}_{\alpha \in A}$ is a family of subsolutions which is bounded from above (w.r.t. the partial ordering on X) then $\sup x^{(\alpha)}$

defined by

$$\left(\sup_{\alpha} x^{(\alpha)} \right)_i = \sup_{\alpha} (x_i^{(\alpha)})$$

is a subsolution.

Analogously, a family of supersolutions $x^{(\alpha)}$ which is bounded from below has an infimum, and $\inf x^{(\alpha)}$ is again a supersolution.

Proof: Part (a) follows immediately from the continuity of the function

Δ .

To prove part (b), let $x^{(\alpha)}$ be a family of subsolutions, indexed by $\alpha \in A$, and let $\bar{x} = \sup x^{(\alpha)}$. Then for each $j \in \mathbb{Z}$ there exists a sequence of α_{jk} 's for which

$$x_j^{(\alpha_{jk})} > \bar{x}_j - 2^{-k} \quad (k = 1, 2, 3, \dots, j \in \mathbb{Z})$$

holds. Clearly $\bar{x} = \sup_{j,k} x_j^{(\alpha_{jk})}$ so that we may assume that the family of $x^{(\alpha)}$'s is countable. Moreover, if one defines

$$x^{(N)} = \sup \{ x_j^{(\alpha_{jk})} \mid 0 \leq k, |j| \leq N \}$$

then, as $N \uparrow \infty$, the $x^{(N)}$ converge in the product topology on X to \bar{x} .

In view of part (a) of this lemma we only have to consider the case in which the number of $x^{(\alpha)}$'s is finite. In turn, this may be reduced to the case of only two $x^{(\alpha)}$'s, by means of an induction argument.

So consider two subsolutions $x, y \in X$ and let $\bar{x} = \sup(x, y)$ be their maximum or supremum. Let $j \in \mathbb{Z}$ be given, and assume that $x_j > y_j$. Then for any nonzero integer k in the range $-l < k < +m$ one has $\bar{x}_{j+k} > x_{j+k}$, so that the monotonicity of Δ (i.e. hypothesis 2.5^a) implies that

$$\Delta(\bar{x}_{j-l}, \dots, \bar{x}_{j+m}) \geq \Delta(x_{j-l}, \dots, x_{j+m}) \geq 0.$$

If $x_j \leq y_j$ then one has, for the same reasons,

$$\Delta(\bar{x}_{j-l}, \dots, \bar{x}_{j+m}) > \Delta(y_{j-l}, \dots, y_{j+m}) > 0 .$$

In either case we obtain $\Delta(\bar{x}_{j-l}, \dots, \bar{x}_{j+m}) > 0$ for arbitrary j , so that \bar{x} is a subsolution. This finishes the proof of the lemma.

The next result shows how subsolutions and supersolutions may be used to construct actual solutions of the recurrence relation (2.4).

Theorem 4.2. Let $\underline{x}, \bar{x} \in X$ be a sub- and a supersolution, respectively which are ordered: $\underline{x} \leq \bar{x}$. Then there is at least one solution of (2.4), say x , between \underline{x} and \bar{x} , i.e. for which $\underline{x} < x < \bar{x}$ holds.

Proof: Define

$$\underline{S} = \{x \in X \mid x \text{ is a subsolution and } x < \bar{x}\} .$$

Clearly \underline{S} is bounded from above, so that

$$x = \sup \underline{S}$$

exists. Since $\underline{x} \in \underline{S}$ we have $\underline{x} < x < \bar{x}$. By our previous lemma x is again a subsolution, and therefore $x \in \underline{S}$. We have for every integer j

$$(4.2) \quad \Delta(x_{j-l}, \dots, x_{j+m}) > 0 .$$

We claim that equality holds for all j , i.e. that x is a solution of (2.4).

To reach a contradiction assume that for some j strict inequality holds in (4.2). Then we must have $x_j < \bar{x}_j$, or $x_j = \bar{x}_j$.

In the first case we define

$$x_k^\epsilon = \begin{cases} x_j + \epsilon & \text{if } j = k \\ x_j & \text{if } j \neq k \end{cases} .$$

The monotonicity of Δ implies that $\Delta(x_{j-l}^\epsilon, \dots, x_{j+m}^\epsilon) > 0$ whenever $k \neq j$ and $\epsilon > 0$. Since we have strict inequality in (4.2) and Δ is continuous we also have $\Delta(x_{j-l}^\epsilon, \dots, x_{j+m}^\epsilon) > 0$ if $\epsilon > 0$ is small enough. So for small $\epsilon > 0$, x^ϵ belongs to \underline{S} , but $x^\epsilon > x$ which is a contradiction.

In the other case, i.e. $x_j = \bar{x}_j$, we observe that, due to the monotonicity of Δ ,

$$\Delta(\bar{x}_{j-l}, \dots, \bar{x}_{j+m}) \geq \Delta(x_{j-l}, \dots, x_{j+m}) > 0.$$

But \bar{x} is a supersolution so that we have a contradiction again.

We must conclude therefore that $x = \sup \underline{S}$ is a solution of the recurrence relation.

It should be clear from the proof that if we had defined

$$\bar{S} = \{x \in X \mid x > \underline{x} \text{ and } x \text{ is a supersolution}\}$$

then $\inf \bar{S}$ would also be a solution of the recurrence relation. In general one expects $\inf \bar{S}$ and $\sup \underline{S}$ to be different.

The theorem can also be used to construct Birkhoff orbits or sequences.

Addendum (4.3). If at least one of \underline{x} and \bar{x} is a Birkhoff sequence,
then there exists a solution x of (2.4) which lies between \underline{x} and \bar{x} , and
is a Birkhoff sequence.

Proof: Let \bar{x} be a Birkhoff sequence. Then we shall prove that $x = \sup(\underline{S})$ also is a Birkhoff sequence.

Let m, n be given integers. Then we have $\tau_{m,n}(\bar{x}) < \bar{x}$ or $\tau_{m,n}(\bar{x}) > \bar{x}$.

In the first case we have, for any $y \in \underline{S}$,

$$\tau_{m,n}(y) \leq \tau_{m,n}(\bar{x}) \leq \bar{x}$$

so that $\tau_{m,n}(y) \in \underline{S}$ (note that the translations $\tau_{m,n}$ preserve the ordering on X). Thus $\tau_{m,n}(\underline{S}) \subset \underline{S}$ and

$$\tau_{m,n}(x) = \tau_{m,n}(\sup \underline{S}) = \sup \tau_{m,n}(\underline{S}) < \sup \underline{S} = x.$$

In the other case we have $\tau_{-m,-n}(\bar{x}) \leq \bar{x}$, so that the same argument shows that $\tau_{-m,-n}(x) < x$ or, $\tau_{m,n}(x) > x$.

The conclusion is, that for any $m, n \in \mathbb{Z}$, one has either $\tau_{m,n}(x) < x$

or $\tau_{m,n}(x) > x$, so that x is a Birkhoff sequence.

If \underline{x} is not a Birkhoff sequence, but \bar{x} is then one shows in the same way that $\inf(\bar{S})$ is a Birkhoff sequence.

Addendum (1.4). If at least one of \bar{x} or x is periodic of type (p,q) , then there exists a solution x of (2.4) which lies between x and \bar{x} , and is periodic of type (p,q) .

Proof: Let \bar{x} be periodic of type (p,q) . Then $\tau_{p,q}(\bar{x}) = \bar{x}$, and therefore $y \in \underline{S}$ implies $\tau_{p,q}(y) \in \underline{S}$. But then $\tau_{p,q}(\sup \underline{S}) = \sup \underline{S}$, so that $\sup \underline{S}$ is periodic of type p,q .

Likewise, if \underline{x} is periodic of type (p,q) then so is $\inf \bar{S}$.

§5. A generalization of Hall's theorem.

In [Ha 1] G. R. Hall proved that any monotone twist map, which has a periodic orbit of type (p,q) , must have a Birkhoff orbit of the same type. The following theorem generalizes this result.

Theorem (5.1). Let $x \in X$ be a solution of the monotone recurrence relation (2.4) for which one can find a real number ω such that

$$M = \sup_{k \in \mathbb{Z}} |x_k - k \cdot \omega| < \infty.$$

Then (2.4) has a Birkhoff solution whose rotation number equals ω .

Proof: Any translate $\tau_{m,n}(x)$ of x is also a solution. Our assumption implies that

$$\underline{x} = \sup\{\tau_{m,n}(x) \mid n \leq m \cdot \omega\}$$

and

$$\bar{x} = \inf\{\tau_{m,n}(x) \mid n > m \cdot \omega\}$$

both exist and that

$$0 < \underline{x}_k - x_k = \sup_{n < m\omega} (x_{k-m} + n - x_k) \leq 2M$$

$$0 < x_k - \bar{x}_k \leq 2M$$

holds. So if we define $\underline{y}_k = \underline{x}_k - 2M$ and $\bar{y}_k = \bar{x}_k + 2M$ then \underline{y} is a subsolution, \bar{y} is a supersolution and $\underline{y} \leq \bar{y}$.

Moreover, \underline{y} and \bar{y} are Birkhoff sequences.

To see this let $r, s \in \mathbb{Z}$ be given and consider $\tau_{r,s}(\underline{x})$:

$$\begin{aligned} \tau_{r,s}(\underline{x}) &= \sup(\tau_{m+r, n+s}(\underline{x}) \mid n < m\omega) \\ &= \sup(\tau_{m,n}(\underline{x}) \mid n < m\omega + s - r\omega) \\ &\geq \underline{x} \quad \text{if } s > r\omega \\ &\leq \underline{x} \quad \text{if } s < r\omega. \end{aligned}$$

Thus \underline{x} is a Birkhoff sequence, and therefore \underline{y} is one too. A similar argument proves that \bar{y} is a Birkhoff sequence.

The addendum to theorem 4.2 implies that the recurrence equation (2.4) has a Birkhoff solution y , between \underline{y} and \bar{y} . Clearly y must have rotation number ω .

§6. Quasiperiodic orbits with prescribed rotation number.

In this section we assume that for some $\omega_0 < \omega_1$ there exist a subsolution \underline{x} and a supersolution \bar{x} of (2.4) such that

$$\begin{aligned}
(6.1) \quad & \liminf_{k \rightarrow \infty} \frac{\bar{x}_k}{k} > \omega_1 \\
& \limsup_{k \rightarrow -\infty} \frac{\underline{x}_k}{k} > \omega_1 \\
& \liminf_{k \rightarrow -\infty} \frac{\bar{x}_k}{k} \leq \omega_0 \\
& \limsup_{k \rightarrow +\infty} \frac{\underline{x}_k}{k} \leq \omega_0
\end{aligned}$$

holds. The main result of this section is:

Theorem 6.1. For any $\omega \in [\omega_0, \omega_1]$ there is a Birkhoff solution of (2.4) with rotation number ω .

Proof: The inequalities (6.1) imply that

$$\liminf_{k \rightarrow +\infty} |k|^{-1} \cdot (\bar{x}_k - \omega \cdot k) \geq \min(\omega_1 - \omega, \omega - \omega_0) > 0$$

and

$$\limsup_{k \rightarrow -\infty} |k|^{-1} \cdot (\underline{x}_k - \omega \cdot k) \leq \max(\omega - \omega_1, \omega_0 - \omega) < 0.$$

Hence there is an integer $N > 0$ such that for all $k \in \mathbb{Z}$ one has:

$$(6.2) \quad \bar{x}_k > \omega k - N \quad \text{and} \quad \underline{x}_k < \omega k + N.$$

Now define

$$\bar{y} = \inf(\tau_{m,n}(\bar{x}) + N \mid n > m\omega)$$

$$\underline{y} = \sup(\tau_{m,n}(\underline{x}) - N \mid n < m\omega).$$

Then, on the ground of arguments similar to those which were used in the proof of theorem 5.1, one concludes that \bar{y} is a Birkhoff supersolution with rotation number ω , and that \underline{y} is a Birkhoff subsolution with the same rotation number.

Furthermore (6.2) implies that

$$\bar{y}_k > \omega k > \underline{y}_k,$$

i.e. that $\underline{y} \leq \bar{y}$. By theorem 4.2 and the addendum (4.3) there must be a Birkhoff solution y between \underline{y} and \bar{y} . This Birkhoff solution has rotation number ω .

§7. A criterion for positive entropy.

Under the hypotheses of section 6 we can construct a great number of orbits of φ_Δ . This construction leads us to the following conclusion.

Theorem 7.1. Let the monotone recurrence relation (2.4) have a sub- and a supersolution, \underline{x} and \bar{x} respectively, which satisfy (6.1) for some $\omega_0 < \omega_1$.


Then the map φ_Δ has positive topological entropy. In fact there is a compact subset K of A^{l+m} such that some iterate of φ_Δ leaves K invariant, and has a Bernoulli shift as a factor, when restricted to K .

Since the existence of K implies positivity of the topological entropy of φ_Δ we only have to construct it to prove the theorem.


The construction proceeds in three steps. In the first step we use \underline{x} and \bar{x} to construct a nicer pair of sub and supersolutions \underline{w} and \bar{w} , which are wedge shaped, as in figure 1.

Then, in the next step we consider a biinfinite sequence of translates of \underline{w} and \bar{w} . These translates of our original wedges are chosen in such a manner that, if \underline{W} denotes the supremum of all subsolutions in question, and \bar{W} the infimum of the translated supersolutions, then $\underline{W} \leq \bar{W}$ and from the results in section four we know that there is a solution W between \underline{W} and \bar{W} . It turns out that we have so much freedom in choosing the translated sub- and supersolutions that we can let the constructed solution W follow any "zig-zag pattern" we like (see figure 2).

Figure 1. Graphs of \underline{w} and \overline{w} .



Figures 2.



Now let's fill in the details.

The wedge like sub and supersolution

Choose two rational numbers ρ_0, ρ_1 such that

$$\omega_0 < \rho_0 < \rho_1 < \omega$$

holds, and define the functions

$$z(t) = \rho_0 t_+ - \rho_1 t_-$$

$$Z(t) = \rho_1 t_+ - \rho_0 t_-$$

where $t_+ = \max(t, 0)$ and $t_- = \max(-t, 0)$.

From our sub- and supersolution \underline{x} and \bar{x} we construct new sub and supersolutions

$$(7.1) \quad \begin{aligned} \underline{w} &= \sup\{\tau_{m,n}(\underline{x}) : n < z(m)\} \\ \bar{w} &= \inf\{\tau_{m,n}(\bar{x}) : n > Z(m)\} \end{aligned}$$

Lemma 4.1 assures us that \underline{w} is a subsolution, and that \bar{w} is a supersolution.

Our hypothesis (6.1) on \underline{x} and \bar{x} implies that

$$\bar{x}_k > z(k) - M$$

and

$$\underline{x}_k < z(k) + M$$

holds for all $k \in \mathbb{Z}$, with M independent of k . For any (m, n) with $n \leq z(m)$ one has

$$\begin{aligned} (\tau_{m,n} \underline{x})_k &= \underline{x}_{k-m} + n \\ &\leq z(k-m) + M + z(m) \\ &\leq z(k) + M \end{aligned}$$

(note that $-z(t)$ is sub additive, so that $-z(k) > -z(k-m) - z(m)$). Thus after taking the supremum over all m, n with $n < z(m)$ one finds

$$\underline{w}_k \leq z(k) + M .$$

Furthermore, if one puts $m = k$ and $n = \lfloor z(k) \rfloor$ (the largest integer below $z(k)$) then

$$\underline{w}_k \geq (\tau_{m,n} \underline{x})_k = \underline{x}_0 + \lfloor z(k) \rfloor .$$

Without loss of generality we may assume that the constant M is so large that $M > |\underline{x}_0| + 1$. Then we have just proved that

$$(7.2) \quad |\underline{w}_k - z(k)| \leq M$$

for all $k \in \mathbb{Z}$.

A similar argument gives

$$(7.3) \quad |\overline{w}_k - z(k)| \leq M$$

(it may be necessary to increase the constant M one more time).

Construction of a zig-zag solution

In the previous step we had chosen two rational numbers ρ_0, ρ_1 . Let Q be a common multiple of their denominators, so that $\rho_0 Q$ and $\rho_1 Q$ are integers. We choose Q so large that

$$(7.4) \quad (\rho_1 - \rho_0) \cdot Q > 4M .$$

Let $\{e_k; k \in \mathbb{Z}\} = e$ be an arbitrary biinfinite sequence of zeroes and ones (i.e. $e_k = 0$ or $e_k = 1$ for all $k \in \mathbb{Z}$). Given such a sequence we define two functions, $\chi_e(t)$ and $\zeta_e(t)$. On the interval $jQ \leq t < (j+1)Q$ we define

$$\chi_e(t) = \rho_{e_j} ,$$

i.e. ρ_1 if $e_j = 1$ and ρ_0 if $e_j = 0$. The other function is given by

$$\zeta_e(t) = \int_0^t \chi_e(s) ds .$$

For any integer j , $\zeta_e(jQ)$ is an integer, and we always have the inequalities

$$z(t) < z_e(t) < Z(t) ,$$

whatever e is.

Now define

$$\underline{w}_e = \sup\{\tau_{jQ, z_e(jQ)}(\underline{w}) \mid j \in \mathbb{Z}\}$$

$$\bar{w}_e = \inf\{\tau_{jQ, z_e(jQ)}(\bar{w}) \mid j \in \mathbb{Z}\} .$$

Proposition 7.2.

$$|\underline{w}_{e,k} - z_e(k)| \leq M$$

$$|\bar{w}_{e,k} - z_e(k)| \leq M .$$

Proof: For any $m \in \mathbb{Z}$ we have

$$\begin{aligned} & [\tau_{jQ, z_e(jQ)}(\underline{w})]_m - z_e(m) \\ &= \underline{w}_{m-jQ} + z_e(jQ) - z_e(m) \\ &= \underline{w}_{m-jQ} - z(m-jQ) + z(m-jQ) - \int_{jQ}^m \chi_e(s) ds \\ &\leq \underline{w}_{m-jQ} - z(m-jQ) \\ &\leq M \end{aligned}$$

where we have used (7.2) in the last step, and the inequality

$$z(t) \leq \int_x^{x+t} \chi_e(s) ds \quad (\forall t \in \mathbb{R}, x \in \mathbb{R})$$

(which follows from $\rho_0 < \chi_e < \rho_1$) in the second last step. This proves one half of the first inequality. To prove the second half we choose an integer j such that $jQ \leq m < (j+1)Q$.

If $e_j = 0$ then

$$\begin{aligned} \underline{w}_{e,m} &> \tau_{jQ, z_e(jQ)}(\underline{w})_m \\ &= \underline{w}_{m-jQ} + z_e(jQ) \end{aligned}$$

$$> z(m-jQ) + \zeta_e(jQ) - M$$

$$= \rho_0(m-jQ) + \zeta_e(jQ) - M$$

$$= \zeta_e(jQ) - M.$$

In case $e_j = 1$ one compares $\underline{w}_{e,m}$ with $\tau(j+1)Q, \zeta_e((j+1)Q) (\underline{w})_m$ to arrive at the same conclusion.

Therefore the first inequality in the Proposition holds. The proof of the second inequality is analogous to that of the first.

It follows immediately from this proposition that $\underline{w}_e - M$ and $\overline{w}_e + M$ are a sub- and supersolution, and the $\underline{w}_e - M < \overline{w}_e + M$, so that there must be at least one solution W of (2.4) which lies between $\underline{w}_e - M$ and $\overline{w}_e + M$.

Construction of the set K

Let Σ_e denote the set of solutions of (2.4) such that $\underline{w}_e - M < W < \overline{w}_e + M$. We have just seen that Σ_e is nonempty.

Let Σ be the union of all Σ_e where e ranges over all possible $\{0,1\}$ -sequences.

To define K , we recall that it has to be a subset of $\mathbb{R}^{l+m}/\mathbb{Z}$, where \mathbb{Z} acts on \mathbb{R}^{l+m} via $(x_{-l}, \dots, x_{m-1}) + 1 = (x_{-l}+1, \dots, x_{m-1}+1)$.

Then we put

$$K_e = \{(x_{-l}, \dots, x_{m-1}) \bmod \mathbb{Z} \mid x \in \Sigma_e\}$$

and

$$K = \{(x_{-l}, \dots, x_{m-1}) \bmod \mathbb{Z} \mid x \in \Sigma\}.$$

Proposition 7.3. K and the K_e are compact.

Proof: Since Σ_e is contained in the order interval $[\underline{w}_e - M, \overline{w}_e + M]$ it is precompact in the product topology on X . Since it is closed Σ_e is in fact compact.

Each K_e is the image of Σ_e under the projection map from X to $\mathbb{R}^{l+m}/\mathbb{Z}$, which is defined by

$$\pi(x) = (x_{-l}, \dots, x_{m-1}) \pmod{\mathbb{Z}}$$

for $x \in X$. Thus $K_e = \pi(\Sigma_e)$ is also compact (the projection map is continuous).

We claim that Σ is also compact. To verify this we observe that Σ is contained in the order interval $\{x \in X \mid z(k) - M < x_k < z(k) + M\}$ and therefore is precompact. It remains to show that Σ is closed. Since X is metrizable we only have to check for sequential closedness. So let $\{w^{(m)}\}_{m \geq 1}$ be a sequence in Σ which converges to $w^* \in X$. Then for every $m \geq 1$ there is a $\{0,1\}$ sequence $e^{(m)} = \{e_k^{(m)}\}_{k \geq \mathbb{Z}}$ such that $w^{(m)} \in \Sigma_{e^{(m)}}$. Now, by passing to a subsequence if necessary, we can arrange things so that for any k the sequence $\{e_k^{(m)}\}_{m \geq 1}$ is eventually constant. If one runs through the definition of the \underline{w}_e and \overline{w}_e one more time then one sees that this implies that the $\underline{w}_e^{(m)}$ and $\overline{w}_e^{(m)}$ converge to \underline{w}_e^* and \overline{w}_e^* respectively (where $e_k^* = \lim_{m \rightarrow \infty} e_k^{(m)}$).

It follows that $w^* \in \Sigma_{e^*}$ and thus $w^* \in \Sigma$.

Hence Σ is closed, and even compact.

We may conclude the proof of the proposition by noting that $K = \pi(\Sigma)$ is the continuous image of a compact set, and therefore also compact.

Proposition 7.4. The sets K_e are pair wise disjoint.

Proof: Let $(x_{-l}, \dots, x_{m-1}) \in \mathbb{R}^{l+m}$ represent a point in K_e . Then we can form its orbit under the diffeomorphism F_Δ (defined in section 2 - it is a lift of φ_Δ), which gives us a biinfinite sequence $\{x_j\}_{j \in \mathbb{Z}} = x$.

On the other hand, there must be an $\tilde{x} \in \Sigma$ such that $(x_{-l}, \dots, x_{m-1}) \pmod{\mathbb{Z}} = \pi(\tilde{x})$. Since \tilde{x} is a solution of (2.4) it is uniquely determined by its components $(\tilde{x}_{-l}, \dots, \tilde{x}_{m-1})$, and we see that $\tilde{x} = x \pmod{\mathbb{Z}}$.

By proposition 7.2 we see that

$$\begin{aligned}
 & |(\tilde{x}_{(j+1)Q} - \tilde{x}_{j \cdot Q}) - \rho_{e_j} \cdot Q| \\
 (7.5) \quad & \leq 2M + |\zeta_e((j+1)Q) - \zeta_e(jQ) - \rho_{e_j} \cdot Q| \\
 & = 2M.
 \end{aligned}$$

We had assumed that $(\rho_1 - \rho_0) \cdot Q > 4M$, so that the inequality cannot be true for both $e_j = 1$ and $e_j = 0$. In other words, given our point in K_e we can find \tilde{x} , and from \tilde{x} we can compute the sequence e . Hence two different K_e 's cannot overlap.

The proof of the last proposition also shows that we have a continuous mapping $\epsilon : K \rightarrow \mathcal{C}$ where $\mathcal{C} = \{0,1\}^{\mathbb{Z}}$ is the space of all $\{0,1\}$ sequences equipped with the product topology (\mathcal{C} is homeomorphic to the Cantor set).

The mapping ϵ is surjective.

On C we have a homeomorphism, called the shift. It acts by $\sigma(e)_k = e_{k+1}$. Using the fact that if $w \in \Sigma_e$, then

$$\tilde{w} = \tau_{-Q, -P}(w) \text{ with } P = -\rho_{e_j} \cdot Q$$

belongs to $\Sigma_{\sigma(e)}$, and also the fact that

$$(\varphi_\Delta)^Q(\pi(w)) = \pi(\tilde{w})$$

one can verify that

- (i) K is invariant under $(\varphi_\Delta)^Q$
- (ii) $\varepsilon \circ (\varphi_\Delta)^Q = \sigma \circ \varepsilon$, i.e. $(\varphi_\Delta)^Q$ maps K_e onto $K_{\sigma(e)}$.

We can restate this by saying that (C, σ) is a factor of $(K, (\varphi_\Delta)^Q)$. By a standard result of ergodic theory this implies that φ_Δ has positive topological entropy (see [Wa]).

We conclude this section with the following observation. If $e \in C$ is a periodic sequence then the corresponding sub and supersolution, \underline{w}_e and \overline{w}_e , are also periodic of type (k, l) for suitable k and l . By the second addendum to theorem 4.2 we know that there must be a (k, l) -periodic solution w_e between \underline{w}_e and \overline{w}_e . Thus we obtain the existence of many periodic orbits of the map φ_Δ .

Indeed, if the sequence $e \in C$ has period k_0 , then $k = k_0 \cdot Q$. It follows that the number of periodic orbits of φ_Δ whose period divides $k_0 \cdot Q$ is at least 2^{k_0} .

§8. Twist maps with zero entropy

Using the criterion for positivity of the topological entropy which we derived in the last section, we now study monotone twist maps of the (two

dimensional) annulus $A = \mathbb{R} \times S^1$, whose entropy vanishes.

Let φ be such a map, and let F be its lift to \mathbb{R}^2 . We assume it satisfies the conditions which were described in section 2. To any point $P \in A$ we associate the sequence $\{x_k(P)\}$ where $x_k(P)$ is the x coordinate of $F^k(P')$ and P' is a lift of P . The sequence $x_k(P)$ depends on the particular lift P' of P which we choose, but the sequence $x_k(P) - x_0(P)$ doesn't.

We define the forward- and backward-rotation numbers of P to be the limits

$$\rho_{\pm}(P) = \lim_{k \rightarrow \pm\infty} \frac{x_k(P) - x_0(P)}{k},$$

if they exist. We shall allow $\rho_{\pm}(P)$ to be $+\infty$ or $-\infty$.

The main result of this section is:

Theorem (8.1). If the map φ has zero topological entropy, then every $P \in A$ has a forward and backward rotation number.

Our other result is:

Theorem (8.2). If the map φ has zero topological entropy and P is a periodic point of type (p,q) with $\gcd(p,q) = 1$, then the orbit of P is a Birkhoff orbit.

This was originally proven by P. Boyland, using Thurston's classification of surface diffeomorphisms.

We begin our proof of these theorems by assuming that we have transformed the problem of finding orbits of the map φ to the problem of finding solutions to the recurrence equation (2.2).

Let $x, y \in X$ be two biinfinite sequences. We shall say that x and y intersect at the integer k if either

$$(y_k - x_k)(y_{k+1} - x_{k+1}) < 0$$

or

$$y_k = x_k \text{ and } (y_{k-1} - x_{k-1})(y_{k+1} - x_{k+1}) < 0$$

holds.

Clearly x and y are ordered if and only if they do not intersect at any integer k . In particular x is a Birkhoff sequence iff it does not intersect any of its translates, $\tau_{m,n}(x)$.

Given a sequence $x \in X$, it can happen that, for some pair of integers (m,n) , with $m > 0$, x and $\tau_{m,n}(x)$ intersect at an infinite number of positive integers k_1, k_2, k_3, \dots . We shall denote the set of (m,n) for which this occurs by $J(x)$. In addition we define the corresponding set of ratios:

$$R(x) = \left\{ \frac{n}{m} \mid (m,n) \in J(x) \right\} \subset \mathbb{Q}.$$

Lemma 8.3. If $R(x)$ is finite, then x has a forward rotation number.

Proof: Define

$$\bar{\rho} = \limsup_{k \rightarrow \infty} \frac{x_k}{k} \text{ and } \underline{\rho} = \liminf_{k \rightarrow \infty} \frac{x_k}{k}.$$

If x does not have a forward rotation number, then $\underline{\rho} < \bar{\rho}$, and there has to be a rational number a/b such that $\underline{\rho} < a/b < \bar{\rho}$. Moreover we can choose a/b in the complement of $R(x)$.

Then x and $\tau_{b,a}(x)$ intersect only at a finite number of positive integers, and there must exist a $k > 0$ such that for all $j > k$ one has

$$x_j > x_{j-b} + a$$

or, for all $j > k$ one has $x_j < x_{j-b} + a$.

In the first case one proves inductively that $x_{k+lb} > x_k + la$, so that $\underline{\rho} > a/b$ which is impossible. The second case also leads to a contradiction.

Thus we have $\underline{\rho} = \bar{\rho}$.

Note that we cannot exclude the possibility that the forward rotation

number, which was shown to exist, is infinite.

Lemma 8.4. If $R(x)$ contains two or more elements, then the map φ has positive topological entropy.

Proof: Let $a/b < c/d$ be two different elements of $R(x)$, and assume that we actually have $(b,a) \in J(x)$ and $(d,c) \in J(x)$. In particular we have $b > 0$ and $d > 0$.

We shall construct a supersolution \bar{y} whose backward rotation number is a/b , and whose forward rotation number is c/d .

Since x and $\tau_{b,a}(x)$ intersect infinitely often at a positive integer there must exist a $j > 0$ such that

$$x_j < \tau_{b,a}(x)_j = x_{j-b} + a$$

$$x_{j+1} > \tau_{b,a}(x)_{j+1} = x_{j+1-b} + a$$

holds. In other words there must exist a $j > 0$ such that $x_i - \tau_{b,a}(x)_i$ increases from negative to positive at j .

For $i < j$ we define \bar{y}_i as follows:

$$\bar{y}_j = x_j, \quad \bar{y}_{j-1} = x_{j-1}, \dots, \bar{y}_{j-b+1} = x_{j-b+1}$$

$$\bar{y}_{j-b} = x_j - a, \quad \bar{y}_{j-b-1} = x_{j-1} - a, \dots, \bar{y}_{j-2b+1} = x_{j-b+1} - a$$

and in general for $t = 0, 1, 2, \dots, b-1$, and $s > 0$:

$$\bar{y}_{j-sb-t} = x_{j-t} - s \cdot a.$$

Then we claim that for $i < j-1$ one has $\Delta(\bar{y}_{i-1}, \bar{y}_i, \bar{y}_{i+1}) < 0$. Since the sequence \bar{y}_i is periodic we only have to verify this for $i = j-1, j-2, \dots, j-b$.

For $j > i > j-b+1$ one has $\bar{y}_i = x_i$, so that

$$\Delta(\bar{y}_{i-1}, \bar{y}_i, \bar{y}_{i+1}) = \Delta(x_{i-1}, x_i, x_{i+1}) = 0 \quad (j-1 \geq i \geq j-b+2).$$

At $i = j-b+1$ one has

$$\Delta(\bar{y}_{i-1}, \bar{y}_i, \bar{y}_{i+1}) = \Delta(x_{j-b+2}, x_{j-b+1}, x_{j-b+1} - a)$$

$$\leq \Delta(x_{j-b+2}, x_{j-b+1}, x_{j-b}) = 0$$

since Δ is increasing in its third argument, and $x_{j-a} < x_{j-b}$. A similar argument can be used to verify the case $i = j-b$.

It remains to define \bar{y}_i for $i > j$. Since x and $\tau_{d,c}(x)$ also intersect at an infinite number of positive integers there must be a $k > 0$ such that

$$x_k > x_{k-d} + c$$

$$x_{k+1} < x_{k-d+1} + c.$$

holds. Moreover we can choose k (much) larger than j . Given this k we define $\bar{y}_i = x_i$ for $j < i < k$, for any $t = 1, 2, \dots, d$ and $s = 0, 1, 2, \dots$ we put

$$\bar{y}_{k+s \cdot d+t} = x_{k+t} + s \cdot a.$$

As above one easily verifies that \bar{y} is a supersolution. Furthermore \bar{y} has a forward and backward rotation number equal to c/d and a/b respectively.

A similar construction can be used to construct a subsolution y with forward rotation number a/b and backward rotation number c/d .

The sub- and supersolution, y and \bar{y} , satisfy the hypothesis (6.1) so that theorem 7.1 tells us that the map φ has positive topological entropy. This completes the proof of lemma 8.4.

Proof of theorem 8.1. If φ has zero topological entropy, then by lemma 8.4, the set $R(x)$ has at most one element, for any given solution x of (2.2). By lemma 8.3 this sequence must have a forward rotation number.

Observe that the reversed sequence $\tilde{x}_k = x_{-k}$ satisfies the recurrence relation

$$\tilde{\Delta}(\tilde{x}_{i-1}, \tilde{x}_i, \tilde{x}_{i+1}) = 0$$

where $\tilde{\Delta}(r, s, t) = \Delta(t, s, r)$. The map corresponding to this (monotone) recurrence relation is conjugate to the inverse of φ and therefore also has

zero topological entropy. Therefore the sequence \tilde{x} has a forward rotation number, and the sequence x has a backward rotation number.

Proof of theorem 8.2. Let $x \in X$ be a sequence which is periodic of type (p,q) . Then for any (m,n) the sequences x and $\tau_{m,n}(x)$ either do not intersect, or they intersect infinitely often (since they are periodic with the same period).

The periodicity also implies that if x and $\tau_{m,n}(x)$ intersect then x and $\tau_{m+q,n+p}(x)$ do too. So if $(m,n) \in J(x)$, then $(m+q,n+p) \in J(x)$ and both n/m and $(n+p)/(m+q)$ belong to $R(x)$. Since we are assuming that φ has zero topological entropy we can apply lemma 8.4 to conclude that $n/m = (n+p)/(m+q)$. But this implies that $n/m = p/q$.

Our other assumption was that $\gcd(p,q) = 1$. Therefore there is an integer $\ell > 1$ such that $(m,n) = (\ell q, \ell p)$. This cannot be true, however, since we then would have $\tau_{m,n}(x) = x$ so that they do not intersect.

We see that the assumption that x and $\tau_{m,n}(x)$ intersect for some (m,n) leads to a contradiction. So x does not intersect any of its translates, which means that it is a Birkhoff sequence.

§9. Existence of Birkhoff sub and supersolutions in general.

We return to the more general monotone recurrence relation (2.4). Given any $\omega \in \mathbb{R}$ the equation (2.4) need not have a Birkhoff solution with rotation number ω . Instead of this, we have the following.

Theorem 9.1. For any $\omega \in \mathbb{R}$ there is a Birkhoff sequence $x \in B$ such that for some $\lambda \in \mathbb{R}$

$$(9.1) \quad \Delta(x_{k-\ell}, \dots, x_{k+m}) = \lambda \quad (k \in \mathbb{Z})$$

(where λ does not depend on k), and such that x has rotation number ω .

In particular, x is either a subsolution or a supersolution.

Using this theorem we shall also prove the next result.

Theorem 9.2. If (2.4) has a Birkhoff subsolution $x^{(0)}$ and a Birkhoff supersolution $x^{(1)}$ then it also has a Birkhoff solution x .

If ω_j is the rotation number of $x^{(j)}$ then the rotation number of x , ω , can be chosen between ω_0 and ω_1 (i.e., $\omega_0 < \omega < \omega_1$ or $\omega_1 < \omega < \omega_0$, depending on how ω_0 and ω_1 are ordered).

Although there is a superficial resemblance between this theorem and theorem 4.2, they are really different. The difference is of course that we do not assume that the sub and supersolution in the last theorem are in any way ordered.

We would also like to point out that, in the case of monotone twist maps, relation 9.1 reduces to

$$Y(x_{k+1}, x_k) = \bar{Y}(x_{k-1}, x_k) + \lambda$$

in the notation of section two. Therefore solutions of 9.1 correspond to orbits of the "translated twist map" given by

$$F_\lambda(x, y) = (f(x, y), g(x, y) + \lambda) ,$$

(again we use the notation of section two).

These maps have been studied before, e.g. by Chenciner in [Ch, section 6]. Our main motivation for studying (9.1), or the F_λ 's is that they seem to be a natural one-parameter family of maps, or recurrence relations associated to Δ . Moreover theorem 9.1 leads to a convenient proof of theorem 9.2.

We begin the proof of theorem 9.1 with the following observation. If there is a dense set of ω 's in \mathbb{R} , for which the theorem holds then it is true for any real ω . Indeed, suppose we can construct Birkhoff orbits $x^{(j)} \in B$ with rotation numbers $\omega^{(j)}$ and suppose the $\omega^{(j)}$ converge to some

$\omega \in \mathbb{R}$. Then we may assume that the $x^{(j)}$ satisfy $0 < x_0^{(j)} < 1$, and by the compactness criterion of section three (see 3.1) we can extract a convergent subsequence. The limit of this subsequence will be the desired Birkhoff orbit with rotation number ω . (This trick is not without precedent, see [Ka 1].)

The theorem is also true if it holds for any Δ which satisfies the following strict monotonicity condition

$$(9.2) \quad \Delta = \Delta(x_{-l}, \dots, x_{+m}) \text{ is strictly monotone} \\ \text{in each } x_j \text{ except } x_0.$$

To see why this is true, let Δ be any function which satisfies a, b and c from section two. Then

$$\Delta_\epsilon(x_{-l} \dots x_{+m}) = \Delta(x_{-l} \dots x_{+m}) + \epsilon \left(\sum_{-l}^m x_j - (l+m+1)x_0 \right)$$

still satisfies a, b and c, and in addition satisfies condition (9.2), for any $\epsilon > 0$. So for any $\epsilon > 0$, Δ_ϵ has a Birkhoff sequence x^ϵ which satisfies (9.1), and has rotation number ω . Again we may take a cluster point of the x^ϵ as $\epsilon \rightarrow 0$ to obtain a solution of 9.1 for our original Δ with the appropriate rotation number.

So from here on we assume that $\omega = p/q$, and that Δ satisfies (9.2). Moreover we suppose $\gcd(p, q) = 1$, and $q > 4$.

We shall show that a solution of (9.1) must exist by means of a homotopy argument.

Let $B_{p,q}$ denote the set of (p, q) periodic Birkhoff sequences. Then $B_{p,q}$ is not empty; it contains the sequence $x_j = jp/q$. The set $B_{p,q}$ is a closed convex subset of the vectorspace X . If we let $X_{p,q}$ denote the affine subspace of X which comprises all (p, q) -periodic sequences, then $B_{p,q}$ has nonempty interior in $X_{p,q}$.

By the compactness criterion of section three, the set

$$D = \{x \in B_{p,q} : x_0 = 0\}$$

is compact. In fact D is homeomorphic to the $(q-1)$ dimensional unit ball in \mathbb{R}^{q-1} , and the mapping

$$(9.3) \quad x \in B_{p,q} \rightarrow (x_0) \times (0, x_1 - x_0, \dots, x_q - x_0) \in \mathbb{R} \times D$$

is a homeomorphism of $B_{p,q}$ with $\mathbb{R} \times D$.

The \mathbb{Z} -action on $B_{p,q}$ induces a \mathbb{Z} -action on $\mathbb{R} \times D$. This induced action is given by "addition to the first coordinate", i.e. $(x, d) \cdot k \rightarrow (x+k, d)$ if $x \in \mathbb{R}$ and $d \in D$.

We let K denote the quotient

$$K = B_{p,q}/\mathbb{Z}$$

and let ∂K be its boundary $\partial K = (\partial B_{p,q})/\mathbb{Z}$, where $\partial B_{p,q}$ is the boundary of $B_{p,q}$ in $X_{p,q}$.

Using the homeomorphism (9.3) we see that $K \cong S^1 \times D$ and $\partial K \cong S^1 \times S^{q-2}$.

On K we define a continuous mapping into \mathbb{R}^q , given by

$$\delta : K \rightarrow \mathbb{R}^q$$

$$\delta(x)_i = \Delta(x_{i-1}, \dots, x_{i+m}) \quad (1 \leq i \leq q) \quad .$$

At first sight it looks as if δ is only defined on $B_{p,q}$, but the periodicity of Δ makes that δ is also well defined on the quotient.

Given δ we define one more mapping, ϵ :

$$\epsilon_k(x) = \delta_k(x) - \frac{1}{q} \sum_{j=1}^q \delta_j(x) \quad (k = 1, 2, \dots, q) \quad .$$

Thus, ϵ is a map from K to the subspace

$$\Lambda = \{(\delta_1, \dots, \delta_q) \in \mathbb{R}^q \mid \delta_1 + \delta_2 + \dots + \delta_q = 0\}$$

of \mathbb{R}^q . We have constructed ϵ in such a manner that solutions of (9.1) are exactly the zeroes of ϵ .

Lemma 9.2. ϵ does not vanish on ∂K .

Proof. Let x and y be any pair of solutions of (9.1), and assume that $x < y$. Then we claim that the strict monotonicity hypothesis (9.2) implies $x \ll y$.

Indeed, if for some k we would have $x_k = y_k$ then

$$\Delta(x_{k-l}, \dots, x_{k+m}) = \Delta(y_{k-l}, \dots, y_{k+m}) (= \lambda)$$

combined with the strict monotonicity of Δ shows that we really have $x_{k-j} = y_{k-j}$ for $-l < j < +m$. By repeating this argument one finds that $x_i = y_i$, or $x_i < y_i$ holds for all $i \in \mathbb{Z}$.

An $x \in X_{p,q}$ belongs to $B_{p,q}$ if and only if for any pair of integers (k,l) one has $\tau_{k,l}(x) > x$ or $\tau_{k,l}(x) < x$. The same point will belong to the boundary of $B_{p,q}$ if, in addition, for some (k,l) one has $\tau_{k,l}(x) > x$ (or $\tau_{k,l}(x) < x$) but not $\tau_{k,l}(x) \gg x$ (or $\tau_{k,l}(x) \ll x$ respectively).

If this point were a solution of (9.1), then $\tau_{k,l}(x)$ would be one too so that $\tau_{k,l}(x) > x$ would imply $\tau_{k,l}(x) \gg x$. But that would contradict $x \in \partial B_{p,q}$.

We conclude that ε does not vanish on the boundary of K .

This lemma shows that ε defines a mapping from ∂K into $\Lambda \setminus \{0\}$. We consider the associated homomorphism $\pi_{q-2}(\varepsilon) : \pi_{q-2}(\partial K) \rightarrow \pi_{q-2}(\Lambda \setminus \{0\})$ of homotopy-groups. Note that these groups are well defined since we have assumed that $q > 4$.

We already saw that $\partial K \cong S^1 \times S^{q-2}$ so that

$$\pi_{q-2}(\partial K) \cong \pi_{q-2}(S^1) \times \pi_{q-2}(S^{q-2}) \cong \mathbb{Z}.$$

Furthermore, $\Lambda \setminus \{0\}$ has the homotopy type of S^{q-2} so that

$\pi_{q-2}(\Lambda \setminus \{0\}) \cong \mathbb{Z}$. It follows that the homomorphism $\pi_{q-2}(\varepsilon)$ is represented by an integer, which we shall call its degree. This degree is defined up to its sign, which depends on the choice of generators of the groups $\pi_{q-2}(\partial K)$ and

$\pi_{q-2}(\Lambda \setminus \{0\})$ which we have implicitly made.

If the degree of ε is nonzero then ε must vanish somewhere in K . Indeed, if $\varepsilon(x) \neq 0$ for all $x \in K$, then we may write $\varepsilon|_{\partial K} = \varepsilon \circ j$, where $j : \partial K \rightarrow K$ is the inclusion mapping. Since $\pi_{q-2}(K) \cong \pi_{q-2}(S^1 \times D) = 0$ (here we use $q > 4$, and the convexity of D) the composition $\pi_{q-2}(\varepsilon) \circ \pi_{q-2}(j)$ vanishes, so that the degree of ε is zero. Therefore we can complete the proof by showing that this degree is nonzero.

With this end in mind we observe that the set

$$\{\Delta \in C^0(\mathbb{R}^{l+m+1}) \mid \Delta \text{ satisfies } \underline{a}, \underline{b}, \underline{c} \text{ of section 2, and also (9.2)}\}$$

is convex and therefore certainly connected. This means that all the maps ε which we have just considered are homotopic on ∂K , and they all have the same degree. We can compute this degree by looking at any particular example that pleases us.

Consider the following choice of Δ :

$$\Delta(x_{-l}, \dots, x_m) = x_{-l} + \dots + x_m - (l+m+1) \cdot x_0.$$

We choose coordinates ξ_1, \dots, ξ_q on $X_{p,q}$ which are defined by

$$x_k = kp/q + \xi_k \quad (1 \leq k \leq q).$$

For $k \leq 0$ or $k > q+1$ we define ξ_k by assuming that $\xi_{k+q} = \xi_k$ holds.

We may identify $X_{p,q}$ with the group ring of $\mathbb{Z}/q\mathbb{Z}$, so that multiplication on $X_{p,q}$ is defined by convolution of the ξ_k -coordinates:

$$(\xi * \eta)_k = \sum_{j=0}^{q-1} \xi_{k-j} \eta_j.$$

The map $\delta : X_{p,q} \rightarrow \mathbb{R}^q$ is simply given by

$$\delta(\xi) = \gamma * \xi$$

where $\gamma = \sum_{j=-l}^{+m} e_j - (l+m+1) \cdot e_0$, and e_j represents $j \in \mathbb{Z}/q\mathbb{Z}$ in the group

ring of $\mathbb{Z}/q\mathbb{Z}$. The map ε is given by

$$\varepsilon(\xi) = \gamma^* \xi - \text{Av}(\gamma^* \xi) = \gamma^*(\xi - \text{Av}(\xi))$$

where $\text{Av } \xi = (\xi_1 + \dots + \xi_q)/q$.

The characters of $\mathbb{Z}/q\mathbb{Z}$ are given by

$$x_j(e_k) = e^{2\pi i j k / q}$$

for $j, k \in \mathbb{Z}/q\mathbb{Z}$.

One finds that

$$x_j(\gamma) = \sum_{k=-l}^{+m} e^{2\pi i j k / q} = (l+m+1)$$

so that if $j \not\equiv 0 \pmod{q}$ one has

$$|x_j(\gamma) + l+m+1| < l+m+1$$

and thus $\text{Re}(x_j(\gamma)) < 0$, and in particular one sees that $x_j(\gamma) \neq 0$.

This implies that the only zeroes of ε are given by $\xi_k = \text{constant}$, i.e. by $x_k = k \cdot \frac{p}{q} + x_0$. This discussion also allows us to compute the degree of ε . Indeed, the map ε restricted to the subspace of $X_{p,q}$, which is defined by $x_0 = 0$, is linear. Its kernel consists of those ξ for which $x_0(\xi - \text{Av}(\xi)) = 0$, i.e. which are constant. So restricted to $\{x \in X_{p,q} : x_0 = 0\}$ it is injective. It follows that the degree of $\varepsilon|_{\partial D \rightarrow \Lambda \setminus \{0\}}$ is ± 1 . Since the inclusion of ∂D in $\partial K \simeq S^1 \times \partial D$ induces an isomorphism on the $(q-2)$ -dimensional homotopy groups the degree of $\varepsilon : \partial K \rightarrow \Lambda \setminus \{0\}$ is also given by $+1$ or -1 .

The proof of theorem 9.1 is complete.

We turn to theorem 9.2.

Define the following two subsets of R .

$$\text{Rot}_+ = \{\omega \in \mathbb{R} \mid \exists x \in B: x \text{ is a supersolution and } x \text{ has rotation number } \omega\}$$

$$\text{Rot}_- = \{\omega \in \mathbb{R} \mid \exists x \in B: x \text{ is a subsolution and } x \text{ has rotation number } \omega\}.$$

It follows directly from theorem 9.1 that the union of Rot_+ and Rot_- is all of \mathbb{R} . Moreover if we define:

$$\text{Rot}_0 = \{\omega \in \mathbb{R} \mid \text{there exists a Birkhoff solution of rotation number } \omega\}$$

then we have $\text{Rot}_0 = \text{Rot}_+ \cap \text{Rot}_-$. Indeed, the inclusions "c" are trivial, and if ω belongs to Rot_+ and Rot_- then there exist a Birkhoff subsolution \underline{x} and a Birkhoff supersolution \bar{x} , both of which have rotation number ω . For a suitable large integer M one will have $\underline{x} - M < \bar{x} + M$ (in view of the inequalities (3.1)) and theorem 4.2 plus the addendum following it tell us that there exists a Birkhoff solution between $\underline{x} - M$ and $\bar{x} + M$.

Hence $\omega \in \text{Rot}_0$.

Using the compactness property described in section three one easily shows that the sets Rot_+ and Rot_- are closed.

The hypothesis of theorem 9.2 is that $\omega_0 \in \text{Rot}_-$ and $\omega_1 \in \text{Rot}_+$. Assume that $\omega_0 < \omega_1$. Then we have just shown that there are two closed sets $A_\pm = [\omega_0, \omega_1] \cap \text{Rot}_\pm$ whose union is the interval $[\omega_0, \omega_1]$, and neither of which is empty. Since the interval is connected the intersection $A_+ \cap A_-$ is nonempty. Therefore there exists a Birkhoff orbit with rotation number $\omega \in [\omega_0, \omega_1]$. The same line of reasoning can be followed when $\omega_1 < \omega_0$, so that we have completed the proof of theorem 9.2.

If we are dealing with the two lowest dimensional cases $\ell = 0, m = 1$ (i.e. degree one circle maps) and $\ell = m = 1$ (i.e. twist maps of a 2-dimensional annulus) then we can improve the previous results.

Theorem 9.3 ($l = 0, m = 1$). If $\varphi : S^1 \rightarrow S^1$ has degree one then it has a Birkhoff orbit.

Proof. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a lift of φ , i.e. $f \in C^0(\mathbb{R}, \mathbb{R})$ and $f(x) + 1 = f(x+1)$. Then $M = \sup |f(x) - x|$ is finite, and one easily verifies that

$$\underline{x}_n = -n \cdot M, \quad \overline{x}_n = +n \cdot M$$

are a sub- and a supersolution for $\Delta(x_0, x_1) = x_1 - f(x_0)$. By theorem (9.2) we know that there exists a Birkhoff solution of $\Delta(x_n, x_{n+1}) = 0$.

In the next Theorem we consider a monotone twist map φ of the two dimensional annulus $S^1 \times \mathbb{R}$.

Theorem 9.4. If there exist $a < b$ such that φ maps the ring $S^1 \times [a, b]$ into itself then φ has a Birkhoff orbit.

Proof. Let $F = (f, g)$ be lift of φ , and define the functions Y, \overline{Y} and Δ as in the beginning of section two. In addition, we consider two auxiliary functions defined by $f_a(x) = f(x, a)$ and $f_b(x) = f(x, b)$.

Then f_a represents a degree-one circle map, and the previous theorem says that there exists a Birkhoff sequence \underline{x}_n for which $\underline{x}_{n+1} = f_a(\underline{x}_n)$ ($n \in \mathbb{Z}$) holds. If one checks the definition of Y, \overline{Y} and Δ then one finds:

$$\begin{aligned} Y(\underline{x}_n, \underline{x}_{n+1}) &= a \\ \overline{Y}(\underline{x}_{n-1}, \underline{x}_n) &= g(x_{n-1}, Y(x_{n-1}, x_n)) > a \\ \implies \Delta(\underline{x}_{n-1}, \underline{x}_n, \underline{x}_{n+1}) &\geq 0. \end{aligned}$$

So \underline{x}_n is a Birkhoff subsolution. By a similar argument a Birkhoff orbit for the map f_b must be a Birkhoff supersolution for Δ .

It follows that the map φ has a Birkhoff orbit.

§10. The symplectic case.

In this final section we draw the reader's attention to a subclass of monotone recurrence relations whose associated map preserves a symplectic structure. Not surprisingly, these recurrence relations are those which can be derived from a variational principle.

Our construction goes like this. Let $h \in C^2(\mathbb{R}^{d+1})$ satisfy

$$(10.1) \quad h_{x_k x_\ell} > \delta \quad (k \neq \ell)$$

for some constant $\delta > 0$. Then we define the formal sum $W(x)$ for $x \in X$ by

$$W(x) = \sum_{j \in \mathbb{Z}} h(x_j, x_{j+1}, \dots, x_{j+d}) .$$

This sum will not converge in general, but if $x, y \in X$ coincide except for a finite number of components, then the a priori formal expression $W(x) - W(y)$ will lead to a finite sum. Hence one can define the derivative of W at an $x \in X$. One finds that:

$$\begin{aligned} \frac{\partial W}{\partial x_j} &= h_d(x_{j-d}, \dots, x_j) + h_{d-1}(x_{j-d+1}, \dots, x_{j+1}) + \dots \\ &\quad \dots + h_0(x_j, \dots, x_{j+d}) \\ &\stackrel{\text{def}}{=} \Delta(x_{j-d}, \dots, x_{j+d}) . \end{aligned}$$

The requirement that W be stationary at $x \in X$ is therefore equivalent to the recurrence relation

$$(10.2) \quad \Delta(x_{j-d}, \dots, x_{j+d}) = 0 \quad (j \in \mathbb{Z}) .$$

If we assume that our original "generating function" h is periodic in the sense that

$$h(x_0+1, \dots, x_d+1) \equiv h(x_0, \dots, x_d) + \phi$$

holds for some constant $\phi \in \mathbb{R}$ then Δ satisfies the periodicity requirement of section 2. From

$$\frac{\partial}{\partial x_1} \Delta(x_{-d}, \dots, x_{+d}) = h_{d-1,d}(x_{-d}, \dots, x_{-1}) + \dots + h_{0,1}(x_0, \dots, x_d)$$

if $0 < i < d$ and a similar expression for $\partial \Delta / \partial x_1$ if $-d < i < 0$ one sees that Δ also satisfies the monotonicity condition, and the coerciveness condition of section two. We conclude that the recurrence relation (10.2) is monotone as we have defined the term.

The recurrence relation 10.2 therefore defines a C^1 diffeomorphism of the $2d$ -dimensional annulus $A^{2d} \sim S^1 \times \mathbb{R}^{2d-1}$. We denote this map by φ_Δ (as in section 2).

We proceed to construct an invariant symplectic form for φ_Δ .

Consider the "partial action" function

$$S(x_1, \dots, x_d, \bar{x}_1, \dots, \bar{x}_d) = \sum_{j=1}^d h(x_j \dots x_d \bar{x}_1 \dots \bar{x}_j) .$$

If $x \in X$ is a solution of (10.2), then

$$S_1 = S(x_{-d+1}, \dots, x_0, x_1, \dots, x_d) + S(x_1 \dots x_d, x_{d+1}, \dots, x_{2d})$$

is stationary with respect to variations of x_1, \dots, x_d . Regarding S_1 as a function of x_1, \dots, x_N the relation $dS_1 = 0$ may be written as

$$(10.3) \quad \sum \bar{Y}_j(x_{-d+1}, \dots, x_d) dx_j = \sum Y_j(x_1, \dots, x_{2d}) dx_j$$

where

$$Y_j(x_1, \dots, x_d, \bar{x}_1, \dots, \bar{x}_d) = \frac{\partial S}{\partial x_j} \quad (1 < j < d)$$

$$\bar{Y}_j(x_1 \dots x_d, \bar{x}_1 \dots \bar{x}_d) = - \frac{\partial S}{\partial \bar{x}_j} \quad (1 < j < d) .$$

It follows from (10.3) that, if we write θ for $Y_1 dx_1 + \dots + Y_d dx_d$ then

$$\theta - \varphi_\Delta^*(\theta) = dS$$

so that φ_Δ preserves the two form $-d\theta = \omega$. The matrix whose (i, j) entry is $\partial Y_j / \partial x_{i+d}$ is lower-triangular, with positive numbers on the diagonal, so

that the 2-form $d\omega = -(dY_1 \wedge dx_1 + \dots + dY_d \wedge dx_d)$ has maximal rank, and is a symplectic form.

A very special case arises if one takes

$$h(x_0, \dots, x_d) = -\frac{1}{2} \sum_{j=1}^d \gamma_j (x_0 - x_j)^2 + \frac{k}{2\pi} \cos 2\pi x_0$$

with $\gamma_1, \dots, \gamma_d > 0$ and $k \in \mathbb{R}$ constants. The associated Δ is

$$\Delta(x_{-d}, \dots, x_d) = \sum_{j=1}^d \gamma_j (x_{-j} - 2x_0 + x_{+j}) + k \sin(2\pi x_0)$$

and the corresponding map φ_Δ generalizes the so called "standard map" (i.e. the case $d = 1, \gamma_1 = 1$).

As an application of the results of section 7 we note that when

$$k > \sum_{j=1}^d j\gamma_j$$

one has a supersolution $\bar{x}_j = \frac{3}{4} + |j|$ and a subsolution $\underline{x}_j = \frac{1}{4} - |j|$ which satisfy (6.1), so that the map φ_Δ has positive entropy.

By analogy with the Aubry-Le Daeron theory [ALD] one could say that the solutions of (10.2) describe the equilibrium states of a bi-infinite chain of particles in which any sequence of $d + 1$ consecutive particles contributes an amount equal to $-h(x_j, \dots, x_{j+d})$ to the total energy of the chain. The number x_j then represents the position of the j -th particle. See fig. 3. From here on we assume that h is truly periodic, i.e. that $\phi = 0$.

As in the Aubry-Le Daeron theory one can show that the function

$$W(x) = \sum_{i=1}^q h(x_i, \dots, x_{i+d})$$

is well defined on $X_{p,q}/\mathbb{Z}$, and in fact is proper on this space. Hence W achieves its maximum on $X_{p,q}$, and for any (p,q) we obtain the existence of a periodic orbit of φ_Δ of type (p,q) .




Fig. 3. A chain of particles with nearest and second nearest neighbour interaction.

Using the positivity of the mixed derivatives one also shows that for $x, y \in X_{p,q}$ one has

$$W(x) + W(y) \leq W(x \vee y) + W(x \wedge y)$$

where \vee and \wedge denote the usual lattice operations on $X_{p,q}$ (one could adapt the proof on page 520 of [Ma 2]). Thus if x maximizes W on $X_{p,q}$ then so does $\tau_{k,l}x$ for any k, l and one finds that

$$2 W(x) \leq W(x \vee \tau_{k,l}(x)) + W(x \wedge \tau_{k,l}(x)) \leq 2 W(x).$$

Hence $x \vee \tau_{k,l}(x)$ and $x \wedge \tau_{k,l}(x)$ also maximize W , and therefore must be solutions of (10.2). Since our Δ satisfies condition (9.2) the proof of lemma 9.2 indicates that we either have $x \vee \tau_{k,l}(x) \gg x$ or $x \wedge \tau_{k,l}(x) = x$; in other words x and $\tau_{k,l}(x)$ are ordered.

The conclusion is that any $x \in X_{p,q}$ which maximizes W is a Birkhoff sequence.

Finally we note that, just as in the Aubry-Le Daeron case, one can obtain Birkhoff orbits with prescribed rotation number by taking limits of similar orbits with rational rotation number.

REFERENCES

- [ALD] S. Aubry and P. Y. Le Daeron, The discrete Frenkel-Kontorova model and its generalizations, *Physica* 8D (1983) p. 381-422.
- [Bo] P. Boyland, Braid types and a topological method of proving positive entropy, preprint.
- [Ch] A. Chenciner, Bifurcations de points fixes elliptiques II, orbites periodiques et ensembles de Cantor invariantes, *Inventiones. Math.* 80 (1985) p. 81-106.
- [Ha 1] G. R. Hall, A topological version of a theorem of Mather on twist maps, *Ergodic Theory and Dynamical Systems*, 4 (1984) p. 585-603.
- [Ha 2] G. R. Hall, A topological version of a theorem of Mather's on shadowing in monotone twist maps, preprint.
- [Ka 1] A. Katok, Some remarks on Birkhoff and Mather twist theorems, *Ergodic Theory and Dynamical Systems*, 2 (1982) p. 185-194.
- [Ka 2] A. Katok, Lyapunov exponents, entropy and periodic orbits for diffeomorphisms, *Publ. Math. de l'I.H.E.S.*, No 51 (1980) p. 137-174.
- [Ma 1] J. N Mather, More Denjoy minimal sets for area-preserving diffeomorphisms, *Comm. Math. Helv.* 60 (1985) p. 508-557.
- [Ma 2] J. N. Mather, Dynamics of Area Preserving Mappings, preprint.
- [Wa] P. Walters, An introduction to ergodic theory, *Grad. Texts in Math.*, Springer-Verlag, New York (1982).

END

DATE

FILMED

7-88

DTIC